Surname	Centre Number	Candidate Number
Other Names		0



#### **GCSE**

C410U20-1





### **WEDNESDAY, 12 JUNE 2019 - MORNING**

# CHEMISTRY – Component 2 Applications in Chemistry

#### **FOUNDATION TIER**

1 hour 15 minutes

	For Examiner's use only		
	Question Maximum Mark Awarded		
	1.	10	
	2.	5	
Section A	3.	9	
	4.	15	
	5.	6	
Section B	6.	15	
	Total	60	

#### **ADDITIONAL MATERIALS**

In addition to this examination paper you will need a:

- calculator and ruler;
- Resource Booklet.

#### **INSTRUCTIONS TO CANDIDATES**

Use black ink or black ball-point pen.

Write your name, centre number and candidate number in the spaces at the top of this page.

Answer all questions.

Write your answers in the spaces provided in this booklet. If you run out of space, use the additional page(s) at the back of the booklet, taking care to number the question(s) correctly.

#### INFORMATION FOR CANDIDATES

This paper is in two sections.

**Section A** (45 marks). You are advised to spend about 50 minutes on this section.

Section B (15 marks). You are advised to spend about 25 minutes on this section.

The number of marks is given in brackets at the end of each question or part-question.


Question 5 is a quality of extended response (QER) question where your writing skills will be assessed.


The Periodic Table is printed on the back cover of this paper and the formulae for some common ions on the inside of the back cover.

#### **SECTION A**

#### Answer all questions.

1. A group of students carried out an investigation into the factors that affect the rusting of iron. They placed identical iron nails into five separate test tubes as shown in the diagram. The nails were left inside each of the tubes for one week.





(a) Complete the table to identify the conditions the nails were exposed to in tubes 4 and 5. The conditions for tubes 1, 2 and 3 have been completed for you. [2]

	Tube 1	Tube 2	Tube 3	Tube 4	Tube <b>5</b>
Air present?	1	X	1		
Water present?	1	X	X		
Salt present?	Х	X	Х		

(b) Suggest why a bung was used in tube 3. [1]


© WJEC CBAC Ltd. (C410U20-1)

Ξха	m	in	er
Ω	nl	v	

(c)	Give the number of the tube that was set up as the control for the investigation. [1		
(d)	State <b>two</b> controlled variables in <b>all five</b> test tubes. Describe how these were controlled [4]		
	Controlled variable	How it was controlled	
(e)			
	The appearance of each of the nail  Tube	Appearance of nail	
		Appearance of nail	
	Tube	Appearance of nail	
	Tube 1	Appearance of nail rusty	
	Tube  1 2	Appearance of nail rusty no rust	
	Tube  1 2 3	Appearance of nail rusty no rust no rust	
	Tube  1 2 3 4 5	Appearance of nail rusty no rust no rust no rust	

2. A group of students carried out an investigation to compare the coloured sugar coatings found on two different brands of sweets – brand **A** and brand **B**. The different colourings are either made from a single dye or a mixture of dyes. The dyes can be separated using chromatography.

The diagram below shows the chromatogram the students produced.



(a)	Use	the information in the chromatogram to answer parts (i)-(iii).
	(i)	Give the colouring that contains the most soluble dye. [1
		Brand
		Colour
	(ii)	Give the colour that is made of exactly the same dyes in both brands of sweets. [1
		Colour
	(iii)	Give the total number of <b>different</b> dyes that are used to make all the colourings.
		Number of different dyes
(b)	the c	$R_{\rm f}$ value of a substance can be used to identify that substance. The $R_{\rm f}$ value of one odyes on the chromatogram is 0.625. Use the equation below to calculate the distance ed by this dye.
	di	istance moved by the dye = $R_{\rm f}$ $ imes$ distance moved by the solvent front
		Distance moved =cm

**3.** (a) Luke and Connor had been investigating tests that are used to identify positive and negative ions in compounds. Some of the results they recorded are shown below.

Put a circle around their **three incorrect** observations.

[3]

#### Flame tests

Metal ion	Flame colour
Li <sup>+</sup>	red
Na <sup>+</sup>	orange
K <sup>+</sup>	blue
Cu <sup>2+</sup>	green

## Adding dilute nitric acid followed by silver nitrate solution

Halide ion	Colour of precipitate
CI <sup>-</sup>	orange
Br <sup>-</sup>	cream
I-	yellow

### Adding sodium hydroxide solution to metal ions in solution

Metal ion solution	Colour of precipitate
Cu <sup>2+</sup>	blue
Fe <sup>2+</sup>	green
Fe <sup>3+</sup>	white

# Adding dilute hydrochloric acid followed by barium chloride solution

lon	Colour of precipitate
SO <sub>4</sub> <sup>2-</sup>	white

(b) In another part of their investigation, Luke and Connor had been given the task of identifying the ions present in a dietary supplement.



They were told that the tablets were made from calcium carbonate and carried out tests to identify both the calcium and carbonate ions in the compound.

(i)	Describe the test they carried out to show that the tablets contain calcium ions.  Give the expected observation.  [2]
(ii)	They tested for carbonate ions by adding dilute hydrochloric acid.  Name the gas that formed if the tablets contain carbonate ions. Describe the test they carried out to identify this gas. Include the result for the test.
	Gas Test and result

© WJEC CBAC Ltd. (C410U20-1) Turn over.

Examiner only

(iii) Each of the tablets has a thin coating. This coating dissolves when the tablet is swallowed, allowing the calcium carbonate to be released into the body. Calcium carbonate is not soluble.

Luke and Connor used a three step method to separate the calcium carbonate powder from its coating and collect a pure sample of calcium carbonate.

Complete the flow diagram. State what is done and why in steps 2 and 3 of the method. [2]

### Step 1

Place the tablet into a beaker of water to dissolve the coating



Step 2




(	Step 3
$\left( \right)$	

# **BLANK PAGE**

© WJEC CBAC Ltd. (C410U20-1) Turn over.

**4.** (a) Anna carried out a titration experiment to find the mass of citric acid in a litre bottle of lemonade.

The diagram shows the apparatus she used to find the volume of lemonade needed to neutralise 25.0 cm<sup>3</sup> of sodium hydroxide solution.




Anna slowly added the lemonade to the sodium hydroxide solution until the indicator changed colour. A trial run was carried out and the titration was then repeated three times. The volume of lemonade added each time was recorded.

Run	Trial	1	2	3
Volume of lemonade (cm <sup>3</sup> )	21.4	19.8	19.9	20.0

<ul><li>(i) Choose your answers to parts I and II from the box</li></ul>	below.
--	--------

	k	ourette	thermometer	pipette	funnel	beaker	
	I.	Name th	e piece of apparatus	labelled A.			[1]
	II.		e piece of apparatus ation.	that would <b>no</b>	<b>t</b> have been	used whilst ca	rrying [1]
(ii)	State		ose of carrying out a	trial run.			[1]
(iii)	State	whether	or not the results are	repeatable. G	ive a reason	for your answe	er. [1]
(iv)			mean volume of lei	monade need	ed to neutra	lise 25.0 cm <sup>3</sup> c	of the [1]
				Mean volun	ne =		cm³
(v)		student's per litre b	final calculations sho ottle.	wed that the I	emonade coi	ntains 1.75g of	citric
	The lacid.		litre bottle of an en	ergy drink stat	es that it cor	ntains 7.00g of	citric
	neutr	alise 25.0	pproximate volume of the sodium bur answer.	of this energy nydroxide solut	drink that w ion used in tl	rould be requir ne experiment.	red to Give [2]
	Volur	ne	cm	3			
	Reas	son					
•••••							
•••••							

(b) Robert used the apparatus shown to investigate the temperature change which occurs when dilute hydrochloric acid reacts with sodium hydroxide solution.




 $25.0\,\mathrm{cm^3}$  of the sodium hydroxide solution was measured into a polystyrene cup. Hydrochloric acid was then added,  $5.0\,\mathrm{cm^3}$  at a time, to the sodium hydroxide solution and the temperature of the mixture recorded each time.

Volume of acid added (cm <sup>3</sup> )	Temperature (°C)
0	21.0
5.0	22.7
10.0	24.0
15.0	25.1
20.0	26.0
25.0	26.5
30.0	26.7
35.0	26.4
40.0	25.6

(i) Plot the temperature against the volume of acid added on the grid below. Draw a suitable line.

[3]



(ii) Use the graph to answer parts I and II.

ı	Find the	overall	temperature	rico	during	tha	avnariment	
Ι.	riiiu iiie	Overall	terriberature	1156	uuriru	เมเษ	experiment	

[1]

°C

II. Find the volume of acid needed to neutralise all the alkali.

[1]

..... cm<sup>3</sup>

(iii)	It is important to reduce heat loss during this experiment.	Examiner only
	Describe how the amount of heat lost was reduced during the experiment. Suggest what else could be done to further reduce heat loss if the experiment were repeated. [2]	
(iv)	One of the risks in this experiment is that the polystyrene cup falls over spilling the reaction mixture. Suggest an improvement to the method that would prevent this from happening. [1]	

5.	Describe how a teacher can demonstrate the reactions of lithium, sodium and potassium in water to show the trend in reactivity in the Group 1 elements.	C
	You should include expected observations and details of any safety precautions that should be taken. [6 QER]	

#### **SECTION B**

Read the article in the Resource Booklet and answer all the questions that follow.

6.	(a)	(a) State <b>three</b> reasons for recycling steel.			
	(b)	(i)	Suggest why some steel car parts may not be directly reused as shown Figure 1.	in [1]	
		(ii)	Use information from lines 4-8 to calculate the mass of carbon dioxide product when one tonne of iron is recycled. Give your answer to <b>three</b> significant figures		
	(c)	(i)	Mass =tonn  Use information from lines 9-11 to calculate the percentage of steel used by the construction industry that goes to landfill each year.		
		(ii)	Percentage =  Use lines 7-8 to calculate the value of the material sent to landfill in 2017.	% [1]	
			Value = £ milli	on	

(d)	(i)	Give the term that is used to describe steel, tool grade steel, stainless steel ar other mixtures of metals.	nd [1]
	(ii)	Suggest why the addition of tungsten makes steel suitable for use in drill tips. [	[1]
	(iii)	Suggest why stainless steel is used to make cutlery.	[1]
(e)	Desc	ribe the trend in worldwide stainless steel production shown in <b>Figure 3</b> . [2	2]
(f)		s 20-23 provide a method of estimating the mass of stainless steel that was produce ears ago.	∍d
		production 20 years ago = production now $\times \frac{35}{100}$	
	For e	example, the mass produced in 2010 was around 35 Mt and that in 1990 was arour	nd
		production in 1990 = 35 $\times \frac{35}{100}$ = 12.25 Mt	
		is the same value to two significant figures as the graph reading so the methodes a good estimate of stainless steel production in 1990.	bd
	Evalu	uate whether this method is suitable for estimating the production in 1970.	3]

Additional page.	Examiner only

### FORMULAE FOR SOME COMMON IONS

POSITIV	E IONS	NEGATI	NEGATIVE IONS		
Name	Formula	Name	Formula		
aluminium	Al <sup>3+</sup>	bromide	Br <sup>-</sup>		
ammonium	$NH_4^+$	carbonate	CO <sub>3</sub> <sup>2-</sup>		
barium	Ba <sup>2+</sup>	chloride	CI <sup>-</sup>		
calcium	Ca <sup>2+</sup>	fluoride	F-		
copper(II)	Cu <sup>2+</sup>	hydroxide	OH <sup>-</sup>		
hydrogen	H⁺	iodide	I-		
iron(II)	Fe <sup>2+</sup>	nitrate	NO <sub>3</sub>		
iron(III)	Fe <sup>3+</sup>	oxide	O <sup>2-</sup>		
lithium	Li⁺	sulfate	SO <sub>4</sub> <sup>2-</sup>		
magnesium	Mg <sup>2+</sup>		·		
nickel	Ni <sup>2+</sup>				
potassium	$K^{+}$				
silver	$Ag^{+}$				
sodium	Na⁺				
zinc	Zn <sup>2+</sup>				

© WJEC CBAC Ltd. (C410U20-1) Turn over.

	7
	9
	2
	4
	က
THE PERIODIC TABLE	Group

0	Helium	20 Neon	40 Argon	84 <b>Kr</b> Krypton 36	Xe Xenon 54	222 Rn Radon 86	
^		19 F Fluorine 9	35.5 Cl Chlorine 17	80 <b>Br</b> Bromine 35	127 	210 At Astatine 85	
9		16 O Oxygen 8	32 S Sulfur 16	79 Se Selenium 34	128 Te Tellurium 52	210 Po Polonium 84	
2		14 Nitrogen 7	31 Phosphorus 15	75 As Arsenic	Sb Antimony 51	209 Bi Bismuth	
4		12 C Carbon 6	28 Silicon	73 <b>Ge</b> Germanium 32	Sn Tin	207 Pb Lead 82	
က		11 B Boron 5	27 AI Aluminium 13	70 <b>Ga</b> 31	115 In Indium 49	204 TI Thallium 81	
				65 Zn Zinc 30	Cd Cadmium 48	201 Hg Mercury 80	
				63.5 Cu Copper 29	Ag Ag Silver	Au Au Gold 79	
				59 <b>Ni</b> Nickel 28	106 Pd Palladium 46	195 <b>Pt</b> Platinum 78	
				59 Co Cobalt 27	103 <b>Rh</b> Rhodium 45	192  r  ridium  77	
Group	e	]		56 Fe Iron 26	101 <b>Ru</b> Ruthenium 44	190 Os Osmium 76	Key
Gro	Hydrogen			55 Mn Manganese 25	99 Tc Technetium 43	186 <b>Re</b> Rhenium 75	
				52 Cr Chromium 24	96 Mo Molybdenum 42	184 W Tungsten 74	
					93 <b>Nb</b> Niobium 41		
				48 Ti Titanium 22	91 Zr Zirconium 40	179 Hf Hafnium 72	
				Scandium 21	89 Y Yttrium 39	139 <b>La</b> Lanthanum 57	227 Actinium 89
7		9 Be Beryllium	24 Mg Magnesium 12	40 Ca Calcium 20	88 Sr Strontium 38	137 Ba Barium 56	226 <b>Ra</b> Radium 88
		I				I	

relative atomic mass

© WJEC CBAC Ltd.

(C410U20-1)

39 **K** Potassium 19

Rubidium 37 133 Cs Caesium 55 Fr Francium 87

Na Sodium